

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

18

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

THIS PAGE BLANK (USPTO)

Ref 2-98

1 HAROLD J. MCELHINNY
ANDREW E. MONACH
2 RACHEL KREVANS
CONSTANCE T.Y. MCCOMB
3 MORRISON & FOERSTER
345 California Street
4 San Francisco, California 94104-2675
Telephone: (415) 677-7000
5
6 Attorneys for Plaintiff
CHIRON CORPORATION

CALENDARED
MORRISON & FOERSTER
MAY 04 1995
FOR DATE(S) _____
BY ES

9 UNITED STATES DISTRICT COURT
10 NORTHERN DISTRICT OF CALIFORNIA

12 CHIRON CORPORATION,
13 Plaintiff,
14 v.
15 ABBOTT LABORATORIES,
16 Defendant.

C 93 4380 MHP

EXPERT REPORT OF B. MATIJA
PETERLIN, M.D. PURSUANT TO
FEDERAL RULE OF CIVIL
PROCEDURE 26

18 I. QUALIFICATIONS

19 I am a Professor of Medicine, Microbiology and Immunology, at the
20 University of California, San Francisco, and an Investigator, Howard Hughes Medical
21 Institute at UCSF. My primary area of research has been focussed for the past ten
22 years on the study of the molecular biology of HIV. A copy of my curriculum vitae is
23 attached hereto as Exhibit A.

1 II. OPINIONS

2 A. Chiron's expression from the EcoRI-KpnI fragment from the
3 ARV2 HIV¹ isolate resulted in an immunoreactive Env polypeptide.

4 Chiron's patent application No. 867,501 describes the identification,
5 cloning and expression of the EcoRI-KpnI fragment from ARV2 in COS cells. We
6 now know that this fragment contains open reading frames for Tat, Rev, Vpu, and Env
7 (Gallo 1988). Staining with an AIDS antiserum revealed cytoplasmic fluorescence only
8 in transfected cells ('501 Application, Sanchez-Pescador 1985). Subsequent reports by
9 many investigators confirmed that this cytoplasmic staining was due to the expression
10 of Env. These investigators also identified Env by electrophoretic mobility and
11 immunoreactivity testing of lysates from these cells (Rekosh 1988, Bird 1990, Kimura
12 1994).

13 Schwartz et al., are the only group which did not detect the expression of
14 Env with a fragment similar to the EcoRI-KpnI fragment from ARV2. In a HeLa cell
15 expression system, they detected only Tat (Schwartz 1990). However, they expressed
16 Env from the same fragment in the rabbit reticulocyte lysate, which is a different
17 expression system. Schwartz et al. acknowledged that other investigators expressed
18 Env with such fragments and were puzzled by their inability to do the same in HeLa
19 cells.

20 Given the data in the Chiron patent and subsequent experiments from
21 many groups, an ordinarily skilled scientist in 1984 or today could conclude confidently
22 that the EcoRI-KpnI fragment from ARV2 expressed Env in COS cells.

23

24

25 1 The virus that causes AIDS is now known as HIV, or human immunodeficiency virus. In
26 1984, when it was first isolated and characterized, it had several names given it by various
27 research groups: LAV (by Dr. Luc Montagnier's group at the French Institut Pasteur); HTLV-
28 III (by Dr. Robert Gallo's group at the National Institutes of Health); and ARV (by Chiron's
collaborator Dr. Jay Levy at the University of California, San Francisco). The name "HIV"
was established in 1986, and is used throughout this report for convenience (Coffin 1986).

1 B. An ordinarily skilled scientist in October 1984 could conclude that
2 the peptide expressed from the EcoRI-KpnI fragment of ARV2 was from Env.

3 In October 1984, Tat, Rev, and Vpu were not known (Sanchez-Pescador
4 1985, Ratner 1985, Wain-Hobson 1985, Muesing 1985). An ordinarily skilled scientist
5 following the example set out in the Chiron application would have expected to express
6 Env and would have done so. By the intense cytoplasmic staining of transfected COS
7 cells which is consistent with the presence of Env, the immunofluorescence test
8 described in the application would have confirmed the expression of Env (Rekosh
9 1988, Bird 1990, Kimura 1994). Tat and Rev proteins have been localized to the
10 nucleus and do not stain in the cytoplasm (Hauber 1987, Ruben 1989, Siomi 1990,
11 Cullen 1988, Felber 1989, Hope 1990). Nuclear staining is easily distinguished from
12 cytoplasmic staining in COS cells, which adhere to and spread out on glass or plastic
13 support (Gluzman 1981). Once the immunofluorescence had confirmed Env, the
14 product of the expression system could have been used by an ordinarily skilled scientist
15 as a recombinant Env immunoassay in 1984. The presence (unknown at the time) of
16 Tat, Rev, and Vpu along with Env would not have impeded the effectiveness of the
17 binding of Env with an AIDS antiserum.

18 In the case of the ARV2, the cytoplasmic staining was due to the
19 expression of Env alone. Vpu is an integral membrane protein, which contains 81
20 amino acids and measures 16 kDa (Cohen 1988, Strebel 1988, Maldarelli 1993).
21 However, not only do many viral strains contain stop codons in *vpu*, but the ARV2
22 sequence as disclosed in the Chiron application contains a stop codon at position 38 in
23 *vpu* (Sanchez-Pescador 1985, Cohen 1988, Myers 1994). Since all reported
24 immunoreactive epitopes of Vpu are encoded by sequences beyond this stop codon
25 (Schneider 1990, Kusk 1993), they would not be included in the protein expressed, and
26 an AIDS serum would not recognize this truncated Vpu peptide from ARV2. Thus,

27

28

1 Chiron scientists did not express immunoreactive Vpu from the EcoRI-KpnI fragment
2 of ARV2 in 1984.

3 An ordinarily skilled scientist working in 1984 with a viral isolate which
4 contained full-length Vpu and Env would have expressed Env along with Tat, Rev, and
5 Vpu (Rekosh 1988, Bird 1990, Kimura 1994). Although it was unnecessary to the
6 recombinant Env immunoassay described in example 9 of the Chiron patent
7 application, if an ordinarily skilled scientist had chosen to perform additional analysis
8 on the product of the EcoRI-KpnI fragment expressed in COS cells, routine
9 electrophoretic mobility and immunoreactivity studies of the expressed proteins would
10 have both confirmed the presence of Env and distinguished Env from the unknown HIV
11 proteins (Tat, Rev, and Vpu). The size of Env as indicated on the sequence of ARV2
12 in the Chiron application combined with the art in 1984 taught an ordinarily skilled
13 scientist at the time that Env would be expressed as a large precursor and would be
14 processed into two large polypeptides. Whereas Env polypeptides measure 160, 120,
15 and 41 kDa, Tat, Rev and Vpu measure only 14, 19, and 16 kDa, respectively (Hauber
16 1987, Cohen 1988, Strebel 1988, Cullen 1988, Ruben 1989, Felber 1989, Hope 1990).
17 In October 1984, as predicted by the ARV2 sequence described in the Chiron
18 application, Tat, Rev, and Vpu would all have been demonstrably too small to be Env.

19 C. Plasmid vectors which could transcribe randomly generated short
20 DNA fragments of unknown nucleotide sequence in all three open reading frames were
21 available for the expression of HIV peptides in 1984 (Gray 1982). However, in
22 October 1984, without accurate sequence information of the fragments and of the
23 original viral genome an ordinarily skilled scientist would not have known from which
24 part of the virus or the host genome these fragments arose.

25 D. In 1984, to generate recombinant polypeptides from Env of any
26 HIV strain, an ordinarily skilled scientist would follow the guidelines of the Chiron
27 patent application. The application states that other isolates of HIV can be found in

28

1 patients with AIDS or ARC. In 1984, methods for obtaining new isolates from lymph
2 nodes or blood of patients were well known in the art (Barre-Sinoussi 1983, Gallo
3 1984, Schupbach 1984, Levy 1984, Luciw 1984). Indeed, the application teaches that
4 two HIV isolates had been identified as of October 1984, one at the Institut Pasteur and
5 the other at the Laboratory of Tumor Cell Biology at The National Institutes of Health
6 (Barre-Sinoussi 1983, Popovic 1984). Two more viral isolates were characterized in
7 the application (ARV3 and ARV4) and their sequences were described as being
8 polymorphic. Thus, the Chiron patent application teaches that different isolates of HIV
9 will have variability in sequence and that more viral isolates should be obtained and
10 characterized.

11 Routine collaborations with hospitals serving communities in which the
12 AIDS virus had been observed, as reported in medical journals and in publications such
13 as Morbidity and Mortality Weekly Reports from the Center for Disease Control,
14 would have and did yield sufficient patients and tissues to identify many new strains of
15 HIV (Myers 1994). The Chiron application describes how to grow the virus in tissue
16 culture cells and how to clone the viral genome. The application teaches that DNA
17 hybridization will yield homologous sequences from new genomic or cDNA libraries.
18 It also directs the sequencing of the viral genome, which was routine in 1984.

19 All HIV1 and HIV2 *env* sequences are sufficiently similar to permit the
20 identification of their *env* ORFs with the HIV *env* probe described in the Chiron patent
21 application (Myers 1994). Moreover, the amount of genetic diversity among *env* genes
22 from various HIV1 and HIV2 isolates would not preclude an ordinarily skilled scientist
23 from extrapolating the Chiron disclosure regarding the location of *env* in ARV2 to all
24 other isolates of HIV1 and HIV2 (Clavel 1986, Guyader 1987, Myers 1994).

25 These steps have been followed since 1984, resulting in the rapid
26 accumulation of viral strains and sequences as well as the identification of HIV2 and
27 SIV (Clavel 1986, Hirsch 1986, Hahn 1987, Charneau 1994, Myers 1994). The

29

1 genomic organization of these lentiviruses, which was first outlined in the Chiron
2 application, was confirmed. In HIV1 and HIV2, the location, size and features of *env*
3 are maintained as are the strategies of viral replication and gene expression (Sanchez-
4 Pescador 1985, Guyader 1987, Hirsch 1987). Therefore, an ordinarily skilled scientist
5 could have followed the teachings of the Chiron application together with the art of
6 1984 to collect a variety of additional HIV1 and HIV2 isolates, to clone them, to
7 sequence them, to identify and characterize their *env* genes, and to make and use
8 recombinant Env immunoassays.

9 E. Dr. Jäck asserts that an inordinate number of disparate peptides
10 from different strains of HIV would have to be examined to obtain smaller peptides to
11 use as immunoreactive reagents against Env. Dr. Jäck does not explain how this
12 assertion relates to the construction of a recombinant Env immunoassay in October
13 1984. If, for example, he means to suggest that his "hundreds of millions" of amino
14 acid sequences must be examined to find short peptides which are broadly cross-
15 reactive, he is wrong. In October 1984, an ordinarily skilled scientist would not have
16 needed to take the approach suggested by Dr. Jäck to determine broadly cross-reactive
17 short peptides. At the time, there were a number of well-known principles for
18 predicting the immunoreactivity of a peptide (Sutcliffe 1983, Singh 1984). For
19 example, instead of randomly expressing peptides from different strains of HIV, a
20 small number of viral isolates would be sequenced and compared.

21 The Chiron application teaches that the three viral isolates from
22 San Francisco were polymorphic, i.e., DNA fragments of different lengths were
23 generated following their digestion with various restriction endonucleases. Following
24 this observation, an ordinarily skilled scientist would have looked for differences in
25 protein sequence between viral isolates in 1984. Routine examination of different Env
26 sequences would have revealed hypervariable, variable, and conserved domains of the
27 protein (Veronese 1985, Starcich 1986, Modrow 1987, Alizon 1987, Myers 1994).

28

1 Conserved sequences would then be examined for their hydrophilicity and
2 hydrophobicity profiles and possible secondary structures (Sutcliffe 1983, Singh 1984).
3 In the art of 1984, immunoreactive oligopeptides of predetermined lengths from
4 conserved sequences were known to interact with specific antibodies (Laver & Air
5 1980, Wiley 1981, Benjamin 1984). Such routine approaches were in fact followed
6 and yielded highly immunoreactive peptides from conserved regions of Env (Gnann
7 8/1987, Gnann 9/1987, Bugge 1990, Bottiger 1990, Krowka 1991, Sprengers 1991,
8 Broliden 1992).

9 Based upon the teachings of the Chiron application and what was known
10 in the art in October 1984, there was no need to screen hundreds of millions of variable
11 sequences. The application identified *env* ORF, demonstrated its expression in cells,
12 and demonstrated its polymorphism. Armed with this information, an ordinarily skilled
13 scientist in 1984 could isolate and characterize new viral strains as well as predict,
14 design and test the immunoreactive peptides from Env of HIV1 and HIV2.

15 F. It would not require extensive experimentation to express Env in
16 host cells not known in 1984.

17 The Chiron application does not require the use of any specific expression
18 system, or purport to claim the invention of expression systems. In October 1984,
19 scientists of ordinary skill were routinely expressing heterologous genes in a wide
20 variety of expression systems, including bacteria, plants, yeast, insect cells, and
21 mammalian cells (e.g., Smith 1983, DiNocera 1983, Chernajovsky 1984, Old &
22 Primrose 1985). After Chiron showed that Env expressed in mammalian cells was
23 immunoreactive with an AIDS serum, an ordinarily skilled scientist would easily have
24 combined the teachings of the patent with any of these expression systems known in
25 1984 or later to subclone the *env* fragment identified by the Chiron application into
26 appropriate vectors and determine levels and fidelities of expression of recombinant
27 proteins.

28

1 Moreover, Dr. Jäck is incorrect in stating that the expression systems he
2 lists were not known in 1984. Most expression systems in use today were either known
3 in October 1984, or are very similar in function to systems known in 1984. In fact,
4 Chiron scientists and others have expressed Env in a variety of host cells and viral
5 systems, including E. coli (Crowl 1985, Schneider 1987, Morikawa 1990), yeast (Barr
6 1987), baculovirus (Hu 1987, Morikawa 1990); CHO cells (Lasky 1986), COS cells
7 (Sanchez-Pescador 1985), vaccinia virus (Chakrabarti 1986), and Drosophila cells
8 (Ivey-Hoyle 1990). WOP cells and expression in sheep were not known in 1984.
9 However, WOP cells represent a system analogous to COS cells (Kern & Basilico
10 1986, Ward 1994), and moreover, neither WOP cells nor expression in sheep has
11 gained wide following in the field of gene expression.

12 G. In October 1984, a scientist of ordinary skill in the field at issue
13 had an advanced degree (a masters, Ph.D., or M.D.) and experience in at least one of
14 the three relevant technical areas: virology, immunology, or molecular biology, as
15 well as some knowledge of immunoassays.

16 H. I have reviewed the opinions expressed by Dr. John Young in his
17 report of May 1, 1995, and I agree with the conclusions expressed therein and the bases
18 therefor.

19 III. COMPENSATION AND PRIOR TESTIMONY

20 I have not testified as an expert at trial or by deposition within the last
21 four years. My compensation in connection with this matter is \$270 per hour.

22 IV. INFORMATION AND REFERENCES CONSIDERED

23 In addition to Dr. Jäck's report and the references listed therein, and my
24 experience in working in the field, as reflected in the publications listed in my
25 curriculum vitae, I considered the content of the following list of articles and I have
26 consulted with Dr. Young regarding the materials set out in Dr. Young's report at
27 Section IV in forming my opinions. To exemplify the basis for the opinions set forth

28

1 above, I have referred to certain of these references in the text. These textual
2 references are meant as examples, not as the exclusive support for each point.

- 3 1. Abacioglu et al., "Epitope mapping and topology of baculovirus-expressed
4 HIV-1 gp160 determined with a panel of murine monoclonal antibodies," AIDS
5 Research and Human Retroviruses 10: 371-381 (1994).
- 6 2. Ahmad et al., "Stable expression of the transfected HIV-1 *env* gene in a
7 human B cell line: characterization of gp120-expressing clones and immunobiological
8 studies," Virology 192: 447-457 (1993).
- 9 3. Alizon et al., "Molecular cloning of lymphadenopathy-associated virus,"
10 Nature 312: 757-760 (12/20-27/84).
- 11 4. Alizon et al., "Genetic variability in human immunodeficiency viruses,"
12 Ann. N.Y. Acad. Sci. 511: 376-384 (1987).
- 13 5. Back et al., "Antibody reactivity to deletion mutants of the HIV-1 SF2
14 envelope," Intervirology 32: 160-172 (1991).
- 15 6. Barr et al., "Antigenicity and immunogenicity of domains of the human
16 immunodeficiency virus (HIV) envelope polypeptide expressed in the yeast
17 *Saccharomyces cerevisiae*," Vaccine 5: 90-101 (6/87).
- 18 7. Barre-Sinoussi et al., "Isolation of a T-lymphotropic retrovirus for a
19 patient at risk for acquired immune deficiency syndrome (AIDS)," Science 228: 868-
20 871 (5/20/83).
- 21 8. Barton et al., "Regeneration of intact tobacco plants containing full length
22 copies of genetically engineered T-DNA, and transmission of T-DNA to R1 progeny,"
23 Cell 32: 1033-1043 (4/83).
- 24 9. Benjamin et al., "The antigenic structure of proteins: a reappraisal,"
25 Ann. Rev. Immunol. 2: 67-101 (1984).
- 26 10. Bird et al., "Expression of human immunodeficiency virus 1 (HIV-1)
27 envelope gene products transcribed from a heterologous promoter," J. Biol. Chem.
28 265: 19151-19157 (11/5/90).
11. Blomberg et al., "A survey of synthetic HIV-1 peptides with natural and
chimeric sequences for differential reactivity with Zimbabwean, Tanzanian and
Swedish HIV-1-positive sera," AIDS 7: 759-767 (6/93).
12. Bottiger et al., "Envelope cross-reactivity between human
immunodeficiency virus types 1 and 2 detected by different serological methods:
correlation between cross-neutralization and reactivity against the main neutralizing
site," J. Virol. 64: 3492-3499 (7/90).
13. Broliden et al., "Specific synthetic peptides for detection of and
discrimination between HIV-1 and HIV-2 infection," J. AIDS 4: 952-958 (1991).

- 1 14. Broliden et al., "Identification of human neutralization-inducing regions of
2 the human immunodeficiency virus type 1 envelope glycoproteins," Proc. Natl. Acad.
3 Sci. USA 89: 461-465 (1/92).
- 4 15. Bugge et al., "Analysis of a highly immunodominant epitope in the human
5 immunodeficiency virus type 1 transmembrane glycoprotein, gp41, defined by a human
6 monoclonal antibody," J. Virol. 64: 4123-4129 (9/90).
- 7 16. Capon et al., "The CD4-gp120 interaction and AIDS pathogenesis," Ann.
8 Rev. Immunol. 9: 649-678 (1991).
- 9 17. Certa et al., "Subregions of a conserved part of the HIV gp41
10 transmembrane protein are differentially recognized by antibodies of infected
11 individuals," EMBO J. 5: 3051-3056 (1986).
- 12 18. Chakrabarti et al., "Expression of the HTLV-III envelope gene by a
13 recombinant vaccinia virus," Nature 320: 535-537 (4/10/86).
- 14 19. Chang et al., "An HTLV-III peptide produced by recombinant DNA is
15 immunoreactive with sera from patients with AIDS," Nature 315: 151-154 (5/9-15/85).
- 16 20. Charneau et al., "Isolation and envelope sequence of a highly divergent
17 HIV-1 isolate: definition of a new HIV-1 group," Virology 205: 247-253 (1994).
- 18 21. Chernajovsky et al., "Efficient constitutive production of human fibroblast
19 interferon by hamster cells transformed with the IFN-beta 1 gene fused to an SV40
20 early promoter," DNA 3: 297-308 (8/84).
- 21 22. Chou et al., "Antibody responses in early human immunodeficiency virus
22 type 1 infection in hemophiliacs," J. Inf. Dis. 157: 805-811 (4/88).
- 23 23. Clavel et al., "Molecular cloning and polymorphism of the human
24 immune deficiency virus type 2," Nature 324: 691-695 (12/18-31/86).
- 25 24. Clerget-Raslain et al., "Specificity of anti-peptide antibodies elicited
26 against synthetic peptides mimicking conserved regions of HIV1 envelope
27 glycoprotein," Res. Virol. 142: 423-438 (1991).
- 28 25. Cloyd & Holt, "Heterogeneity of human immunodeficiency virus cell-
associated antigens and demonstration of virus type specificities of human antibody
responses," Virology 161: 286-292 (1987).
- 26 26. Coffin et al., "What to call the AIDS virus," Nature 321:10 (5/1-7/86).
- 27 27. Cohen et al., "Identification of a protein encoded by the *vpu* gene of
28 HIV-1," Nature 334: 532-534 (8/11/88).
- 29 28. Crowl et al., "HTLV-III *env* gene products synthesized in *E. coli* are
recognized by antibodies present in the sera of AIDS patients," Cell 41: 979-86 (7/85).
- 30 29. Cullen et al., "Subcellular localization of the human immunodeficiency
virus *trans*-acting *art* gene product," J. Virol. 62: 2498-2501 (7/88).

- 1 30. Cumming et al., "Use of conserved immunodominant epitope of HIV
surface glycoprotein gp41 in the detection of early antibodies," AIDS 4: 83-86 (1990).
- 2 31. De et al., "Comparative analysis of nucleotide sequences of the partial
3 envelope gene (5' domain) among human T lymphotropic virus type I (HTLV-I)
isolates," Virology 182: 413-419 (1991).
- 4 32. Dewhurst et al., "Sequence analysis and acute pathogenicity of
5 molecularly cloned SIV smm-pbj14," Nature 345: 636-640 (6/14/90).
- 6 33. de Wolf et al., "Characterization of human antibody-binding sites on the
7 external envelope of human immunodeficiency virus type 2," J. Gen. Virol. 72: 1261-
1267 (1991).
- 8 34. DiNocera et al., "Transient expression of genes introduced into cultured
cells of *Drosophila*," Proc. Natl. Acad. Sci. USA 80: 7095-7098 (12/83).
- 9 35. Dowbenko et al., "Epitope mapping of the human immunodeficiency virus
10 type 1 gp120 with monoclonal antibodies," J. Virol. 62: 4703-4711 (12/88).
- 11 36. DuBois et al., "Expression and purification of protein segments encoded
by the envelope and 3'-orf genes of human immunodeficiency virus type 1," AIDS
12 Res. Hum. Retrovir. 4: 419-431 (1988).
- 13 37. Earl et al., "Folding, interaction with GRP78-BiP, assembly, and
14 transport of the human immunodeficiency virus type 1 envelope protein," J. Virol. 65:
2047-2055 (4/91).
- 15 38. Essex et al., "Antigenic characterization of the human immunodeficiency
16 viruses," J. Am. Acad. Derm. 22: 1206-1210 (6/90).
- 17 39. Felber et al., "Rev protein of human immunodeficiency virus type 1
affects the stability and transport of the viral mRNA," Proc. Natl. Acad. Sci. USA 86:
1495-1499 (3/89).
- 18 40. Franchini et al., "Spectrum of natural antibodies against five HTLV-III
19 antigens in infected individuals: correlation of antibody prevalence with clinical
status," Blood 69: 437-441 (2/87).
- 20 41. Friberg et al., "Mutational analysis of the HIV-1 Vpu protein," Leukemia
21 8 (Suppl. 1): S156-S162 (4/94).
- 22 42. Gallo et al., "Frequent detection and isolation of cytopathic retroviruses
(HTLV-III) from patients with AIDS and at risk for AIDS," Science 224: 500-503
23 (5/4/84).
- 24 43. Gallo et al., "HIV/HTLV nomenclature (letter)," Nature 333: 504
(6/9/88).
- 25 44. Gluzman et al., "SV40-transformed simian cells support the replication of
26 early SV40 mutants," Cell 23: 175-182 (1981).
- 27
- 28

- 1 45. Gnam et al., "Diagnosis of AIDS by using a 12-amino acid peptide
2 representing an immunodominant epitope of the human immunodeficiency virus," J. Inf. Dis. 156: 261-267 (8/87).
- 3 46. Gnam et al., "Synthetic peptide immunoassay distinguishes HIV type 1
4 and HIV type 2 infections," Science 237: 1346-1349 (9/11/87).
- 5 47. Goeddel et al., "Expression in Escherichia coli of chemically synthesized
6 genes for human insulin," Proc. Natl. Acad. Sci. USA 76: 106-110 (1/79).
- 7 48. Gonda, M., "Molecular genetics and structure of the human
8 immunodeficiency virus," J. Elec. Micro. Tech. 8: 17-40 (1988).
- 9 49. Gottlinger et al., "Vpu protein of human immunodeficiency virus type 1
10 enhances the release of capsids produced by gag gene constructs of widely divergent
11 retroviruses," Proc. Natl. Acad. Sci. USA 90: 7381-7385 (8/93).
- 12 50. Gray et al., "Open reading frame cloning: Identification, cloning and
13 expression of open reading frame DNA," Proc. Natl. Acad. Sci. USA 79: 6598-6602
14 (11/82).
- 15 51. Guyader et al., "Genome organization and transactivation of the human
16 immunodeficiency virus type 2," Nature 326: 662-669 (4/16/87).
- 17 52. Hahn et al., "Molecular cloning and characterization of the HTLV-III
18 virus associated with AIDS," Nature 312: 166-169 (11/8/84).
- 19 53. Hahn et al., "Relation of HTLV-4 to simian and human
20 immunodeficiency-associated viruses," Nature 330: 184-186 (11/12/87).
- 21 54. Haigwood, N., "Importance of hypervariable regions of HIV-1 gp120 in
22 the generation of virus neutralizing antibodies," AIDS Res. Hum. Retrovir. 6: 855-869
23 (1990).
- 24 55. Haigwood, N., "Native but not denatured recombinant human
25 immunodeficiency virus type 1 gp120 generates broad-spectrum neutralizing antibodies
26 in baboons," J. Virol. 66: 172-182 (1/92).
- 27 56. Hammariskjold et al., "Regulation of human immunodeficiency virus *env*
28 expression by the *rev* gene product," J. Virol. 63: 1959-1966 (5/89).
57. Hammer et al., "Production of transgenic rabbits, sheep and pigs by
 microinjection," Nature 315: 680-683 (6/20-26/85).
58. Hauber et al., "Trans-activation of human immunodeficiency virus gene
 expression is mediated by nuclear events," Proc. Natl. Acad. Sci. USA 84: 6364-6368
 (9/87).
59. Helseth et al., "Human immunodeficiency virus type 1 gp120 envelope
 glycoprotein regions important for association with the gp41 transmembrane
 glycoprotein," J. Virol. 65: 2119-2123 (4/91).

1 60. Henklein et al., "Synthesis and characterization of the hydrophilic C-
2 terminal domain of the human immunodeficiency virus type 1-encoded virus protein
U (Vpu)," Pent. Res. 6: 79-87 (1993).

3 61. Hirsch et al., "Cross-reactivity to human T-lymphotropic virus type
4 III/lymphadenopathy-associated virus and molecular cloning of simian T-cell
lymphotropic virus type III from African green monkeys," Proc. Natl. Acad. Sci. USA
83: 9754-9758 (12/86).

5 62. Hirsch et al., "The genome organization of STLV-3 is similar to that of
6 the AIDS virus except for a truncated transmembrane protein," Cell 49: 307-319
(5/8/87).

7 63. Ho et al., "Second conserved domain of gp120 is important for HIV
8 infectivity and antibody neutralization," Science 239: 1021-1023 (2/26/88).

9 64. Hope et al., "Steroid-receptor fusion of the human immunodeficiency
10 virus type 1 Rev transactivator: mapping cryptic functions of the arginine-rich motif,"
Proc. Natl. Acad. Sci. USA 87: 7787-7791 (10/90).

11 65. Hu et al., "Expression of envelope glycoproteins of human
12 immunodeficiency virus by an insect virus vector," J. Virol. 61: 3617-3620 (11/87).

13 66. Ivey-Hoyle & Rosenberg, "Rev-dependent expression of human
14 immunodeficiency virus type 1 gp160 in *Drosophila melanogaster* cells," Mol. Cell.
Biol. 10: 6152-6159 (12/90).

15 67. Kern & Basilico, "An inducible eukaryotic host-vector expression system:
16 amplification of genes under the control of the polyoma late promoter in a cell line
producing a thermolabile large T antigen," Gene 43: 237-245 (1986).

17 68. Kimura et al., "Intracellular membrane traffic of human
18 immunodeficiency virus type 1 envelope glycoproteins: Vpu liberates Golgi-targeted
gp160 from CD4-dependent retention in the Endoplasmic Reticulum," J. Biochem.
115: 1010-1020 (1994).

19 69. Kitchen et al., "Viral envelope protein of HTLV-III is the major target
20 antigen for antibodies in hemophiliac patients," J. Inf. Dis. 153: 788-790 (4/86).

21 70. Klasse et al., "A cluster of continuous antigenic structures in the
22 transmembrane protein of HIV-1: individual patterns of reactivity in human sera,"
Mol. Immunol. 28: 613-622 (1991).

23 71. Kleinschmidt et al., "Serological markers as prognostic criteria for the
course of HIV infection," Infection 19 (Suppl. 2): S89-S92 (1991).

24 72. Kornfield et al., "Cloning of HTLV-4 and its relation to simian and
25 human immunodeficiency viruses," Nature 326: 610-613 (4/9/87).

26 73. Kovac et al., "Bacterially expressed core and envelope proteins of human
27 immunodeficiency virus type-1 (HIV-1): comparative evaluation in detection of type-
specific antibodies," Acta. Virol. 36: 337-346 (1992).

- 1 74. Kowalski et al., "Functional regions of the envelope glycoprotein of
human immunodeficiency virus type 1," Science 237: 1351-1355 (9/11/87).
- 2 75. Kozarsky et al., "Glycosylation and processing of the human
3 immunodeficiency virus type 1 envelope protein," J. AIDS 2: 163-169 (1989).
- 4 76. Kreutz et al., "Analysis of the envelope region of the highly divergent
HIV-2alt isolate extends the known range of variability within the primate
5 immunodeficiency viruses," AIDS Res. Hum. Retrovir. 8: 1619-1629 (1992).
- 6 77. Krowka et al., "Epitopes of human immunodeficiency virus type 1 (HIV-
1) envelope glycoproteins recognized by antibodies in the sera of HIV-1-infected
7 individuals," Clin. Immunol. Immunopath. 59: 53-64 (1991).
- 8 78. Kusk et al., "Mapping of a new immunodominant human linear B-cell
epitope on the Vpu protein of the human immunodeficiency virus type 1," J. AIDS 6:
9 334-338 (1993).
- 10 79. Landau et al., "The envelope glycoprotein of the human
immunodeficiency virus binds to the immunoglobulin-like domain of CD4," Nature
11 334: 159-162 (7/14/88).
- 12 80. Lasky et al., "Neutralization of the AIDS retrovirus by antibodies to a
recombinant envelope glycoprotein," Science 233: 209-212 (7/11/86).
- 13 81. Lasky et al., "Delineation of a region of the human immunodeficiency
14 virus type 1 gp120 glycoprotein critical for interaction with the CD4 receptor," Cell
50: 975-985 (9/11/87).
- 15 82. Laver & Air, Structure and Variation in Influenza Virus (Elsevier 1980).
- 16 83. Leonard et al., "Assignment of intrachain disulfide bonds and
17 characterization of potential glycosylation sites of the type 1 recombinant human
immunodeficiency virus envelope glycoprotein (gp120) expressed in Chinese hamster
18 ovary cells," J. Biol. Chem. 265: 10373-10382 (6/25/90).
- 19 84. Levy et al., "Isolation of lymphocytopathic retroviruses from San
Francisco patients with AIDS," Science 225: 840-842 (8/24/84).
- 20 85. Lombardi et al., "Detection of antibodies to human immunodeficiency
21 virus type 1 by using synthetic peptides and time-resolved fluoroimmunoassay (TR-
FIA)," Eur. J. Epidemiol. 8: 298-304 (3/92).
- 22 86. Lomedico et al., "Use of recombinant DNA technology to program
23 eukaryotic cells to synthesize rat proinsulin: a rapid expression assay for cloned genes,"
Proc. Natl. Acad. Sci. USA 79: 5798-5802 (10/82).
- 24 87. Luciw et al., "Molecular cloning of AIDS-associated retrovirus," Nature
25 312: 760-763 (12/27/84).
- 26 88. Luciw et al., "Regulated expression of HIV-1 in CD4+ lymphoid cells,"
Retroviruses of Human AIDS and Related Animal Diseases 62-66 (Girard & Valette,
27 eds., Fondation Marcel Merieux 1987).

- 1 89. Luke et al., "Structural comparison of the external glycoproteins of
human and simian immunodeficiency virus," Intervirology 32: 198-201 (1991).
- 2 90. Mackett et al., "General method for production and selection of infectious
3 vaccinia virus recombinants expressing foreign genes," J. Virol. 49: 857-864 (3/84).
- 4 91. Maldarelli et al., "Human immunodeficiency virus type 1 Vpu protein is
an oligomeric type I integral membrane protein," J. Virol. 67: 5056-5061 (8/93).
- 5 92. McCutchan et al., "Genetic comparison of human immunodeficiency virus
6 (HIV-1) isolates by polymerase chain reaction," J. AIDS 4: 1241-1250 (1991).
- 7 93. Miyajima et al., "Analysis of full-length cDNA clones carrying GAL1 of
8 *Saccharomyces cerevisiae*: a model system for cDNA expression," Nucl. Acids Res.
12: 6397-6414 (8/24/84).
- 9 94. Modrow et al., "Computer-assisted analysis of envelope protein sequences
of seven human immunodeficiency virus isolates: prediction of antigenic epitopes in
10 conserved and variable regions," J. Virol. 61: 570-578 (2/87).
- 11 95. Moore & Jarrett, "Sensitive ELISA for the gp120 and gp160 surface
glycoproteins of HIV-1," AIDS Res. Hum. Retrovir. 4: 369-379 (1988).
- 12 96. Moore et al., "An enzyme-linked immunosorbent assay for antibodies to
13 the envelope glycoproteins of divergent strains of HIV-1," AIDS 3: 155-163 (1989).
- 14 97. Moore et al., "Characterization of recombinant gp120 and gp160 from
HIV-1: binding to monoclonal antibodies and soluble CD4," AIDS 4: 307-315 (1990).
- 15 98. Moore et al., "Monoclonal antibodies to HIV-1 gp120: a request," AIDS
16 Res. Hum. Retrovir. 9: 695 (1993).
- 17 99. Moore, J., "The reactivities of HIV-1 + human sera with solid-phase V3
loop peptides can be poor predictors of their reactivities with V3 loops on native gp120
18 molecules," AIDS Res. Hum. Retrovir. 9: 209-219 (1993).
- 19 100. Moore et al., "Antibodies to discontinuous or conformationally sensitive
epitopes on the gp120 glycoprotein of human immunodeficiency virus type 1 are highly
20 prevalent in sera of infected humans," J. Virol. 67: 863-875 (2/93).
- 21 101. Moore et al., "Immunochemical analysis of the gp120 surface
glycoprotein of human immunodeficiency virus type 1: probing the structure of the C4
22 and V4 domains and the interaction of the C4 domain with the V3 loop," J. Virol. 67:
4785-4796 (8/93).
- 23 102. Moore et al., "Probing the structure of the V2 domain of human
24 immunodeficiency virus type 1 surface glycoprotein gp120 with a panel of eight
monoclonal antibodies: human immune response to the V1 and V2 domains," J. Virol.
25 67: 6136-6151 (10/93).
- 26 103. Moore et al., "Probing the structure of the human immunodeficiency virus
surface glycoprotein gp120 with a panel of monoclonal antibodies," J. Virol. 68: 469-
27 484 (1/94).
- 28

- 1 104. Moore et al., "Exploration of antigenic variation in gp120 from clades A
2 through F of human immunodeficiency virus type 1 by using monoclonal antibodies,"
3 J. Virol. 68: 8350-8364 (12/94).
- 4 105. Morikawa et al., "Expression of HIV-1 gp120 and human soluble CD4 by
5 recombinant baculoviruses and their interaction in vitro," AIDS Res. Hum. Retrovir. 6:
6 765-773 (1990).
- 7 106. Morikawa et al., "HIV-1 envelope protein gp120 expression by secretion
8 in *E. coli*: assessment of CD4 binding and use in epitope mapping," J. Virol. Meth.
9 29: 105-114 (1990).
- 10 107. Muesing, M., "Nucleic acid structure and expression of the human
11 AIDS/lymphadenopathy retrovirus," Nature 313: 450-458 (2/7/85).
- 12 108. Muster et al., "A conserved neutralizing epitope on gp41 of human
13 immunodeficiency virus type 1," J. Virol. 67: 6642-6647 (11/93).
- 14 109. Muster et al., "Cross-neutralizing activity against divergent human
15 immunodeficiency virus type 1 isolates induced by the gp41 sequence ELDKWAS," J.
16 Virol. 68: 4031-4034 (6/94).
- 17 110. Myers et al., Human Retroviruses and AIDS 1994: A Compilation and
18 Analysis of Nucleic Acid and Amino Acid Sequences (Los Alamos Nat'l Lab. 1994).
- 19 111. Norrby et al., "Human immunodeficiency virus antibody responses
20 determined by site-directed serology," Intervirology 31: 315-326 (1990).
- 21 112. Old & Primrose, Principles of Gene Manipulation (Blackwell Scientific
22 Publications, 3rd ed. 1985).
- 23 113. Ou et al., "Wide distribution of two subtypes of HIV-1 in Thailand,"
24 AIDS Res. Hum. Retrovir. 8: 1471-2 (1992).
- 25 114. Peterlin, B., "Transcriptional regulation of HIV-1," Human Retroviruses
26 (UCLA Symposia on Molecular and Cellular Biology) 119: 153-162 (Groopman et al.,
27 eds., Alan R. Liss 1990).
- 28 115. Peterlin, B., "Molecular biology of HIV," Retroviridae (Levy, J., ed.,
Plenum Press 1995).
116. Peterlin et al., "Tat *trans*-activator," Molecular Biology of Human
Retroviruses 75-100 (Cullen, B., ed., ILR/Oxford University Press 1993).
117. Peterlin & Luciw, "Molecular biology of HIV," AIDS 2: S29-S40 (1988).
118. Peterlin & Luciw, "Replication of the human immunodeficiency virus:
strategies for inhibition," Biotechnology 6: 794-799 (7/88).
119. Popovic et al., "Detection, isolation and continuous production of
cytopathic retroviruses (HTLV-III) from patients with AIDS and pre-AIDS," Science
224: 497-500 (5/4/84).

- 1 120. Profy et al., "Epitopes recognized by the neutralizing antibodies of an
2 HIV-1-infected individual," J. Immunol. 144: 4641-4647 (6/15/90).
- 3 121. Ratner et al., "Complete nucleotide sequence of the AIDS virus, HTLV-
4 III," Nature 313: 277-283 (1/24/85).
- 5 122. Re et al., "Patterns of antigenaemia and antibody response in patients
6 infected with human immunodeficiency virus (HIV) according to clinical state," J.
7 Clin. Pathol. 42: 282-283 (1989).
- 8 123. Reed & Kinzel, "Primary structure elements responsible for the
9 conformational switch in the envelope glycoprotein gp120 from human
10 immunodeficiency virus type 1: LPCR is a motif governing folding," Proc. Natl.
11 Acad. Sci. USA 90: 6761-6765 (7/93).
- 12 124. Reiss et al., "Antibody response to viral proteins U (Vpu) and R (Vpr) in
13 HIV-1-infected individuals," J. AIDS 3: 115-122 (1990).
- 14 125. Rekosh et al., "Coexpression of human immunodeficiency virus envelope
15 proteins and Tat from a single simian virus 40 late replacement vector," Proc. Natl.
16 Acad. Sci. USA 85: 334-338 (1/88).
- 17 126. Resnick et al., "Serologic characterization of human immunodeficiency
18 virus infection by Western Blot and radioimmunoprecipitation assays," Arch. Pathol.
19 Lab. Med. 111: 1040-1044 (11/87).
- 20 127. Roben et al., "Recognition properties of a panel of human recombinant
21 Fab fragments to the CD4 binding site of gp120 that show differing abilities to
22 neutralize human immunodeficiency virus type 1," J. Virol. 68: 4821-4828 (8/94).
- 23 128. Robert-Guroff et al., "Cross-neutralization of human immunodeficiency
24 virus type 1 and 2 and simian immunodeficiency virus isolates," J. Virol. 66: 3602-
25 3608 (6/92).
- 26 129. Ruben et al., "Structural and functional characterization of human
27 immunodeficiency virus Tat protein," J. Virol. 63: 1-8 (1/89).
- 28 130. Sanchez-Pescador et al., "Nucleotide sequence and expression of an
AIDS-associated retrovirus (ARV-2)," Science 227: 484-492 (2/1/85).
131. Schneider et al., "A new ELISA test for HIV antibodies using a
bacterially produced viral env gene product," Med. Microbiol. Immunol. 176: 47-51
(1987).
132. Schneider et al., "The antibody response to the HIV-1 specific 'out' (Vpu)
protein: identification of an immunodominant epitope and correlation of antibody
detectability to clinical stages," AIDS Res. Hum. Retrovir. 6: 943-950 (1990).
133. Schneider et al., "Expression of Nef, Vpu, CA and CD4 during the
infection of lymphoid and monocytic cell lines with HIV-1," Arch. Virol. 125: 161-
176 (1992).
134. Schubert et al., "The human immunodeficiency virus type 1 encoded Vpu
protein is phosphorylated by casein kinase-2 (CK-2) at positions Ser52 and Ser56

- 1 within a predicted alpha-Helix-Turn-alpha-Helix-motif," J. Mol. Biol. 236: 16-25
2 (1994).
- 3 135. Schubert et al., "Differential activities of the human immunodeficiency
4 virus type 1-encoded Vpu protein are regulated by phosphorylation and occur in
5 different cellular compartments," J. Virol. 68: 2260-2271 (4/94).
- 6 136. Schulz et al., "Conserved structural features in the interaction between
7 retroviral surface and transmembrane glycoproteins?," AIDS Res. Hum. Retrovir. 8:
8 1571-1580 (1992).
- 9 137. Schupbach et al., "Serological analysis of a subgroup of human T-
10 lymphotropic retroviruses (HTLV-III) associated with AIDS," Science 224: 503-505
11 (5/4/84).
- 12 138. Schwartz et al., "Env and Vpu proteins of human immunodeficiency virus
13 type 1 are produced from multiple bicistronic mRNAs," J. Virol. 64: 5448-5456
14 (11/90).
- 15 139. Schwartz et al., "Mechanism of translation of monocistronic and
16 multicistronic human immunodeficiency virus type 1 mRNAs," Mol. Cell. Biol. 12:
17 207-219 (1/92).
- 18 140. Shoeman et al., "Comparison of recombinant human immunodeficiency
19 virus Gag precursor and Gag/Env fusion proteins and a synthetic Env peptide as
20 diagnostic reagents," Anal. Biochem. 161: 370-379 (1987).
- 21 141. Singh et al., Ind. J. Chem. 23B: 1237 (1984).
- 22 142. Siomi et al., "Effects of a highly basic region of human
23 immunodeficiency virus Tat protein on nucleolar localization," J. Virol. 64: 1803-1807
24 (4/90).
- 25 143. Skinner et al., "Neutralizing antibodies to an immunodominant envelope
26 sequence do not prevent gp120 binding to CD4," J. Virol. 62: 4195-4200 (11/88).
- 27 144. Smith et al., "Production of human beta interferon in insect cells infected
28 with a baculovirus expression vector," Mol. Cell Biol. 3: 2156-2165 (12/83).
145. Smythe et al., "Production of linear polymers of HIV gp120-binding
domains," Protein Engineering 7: 145-147 (1994).
146. Spies et al., "Alternate pathways of secretion of simian immunodeficiency
virus envelope glycoproteins," J. Virol. 67: 6535-6541 (11/93).
147. Sprengers et al., "A microELISA system for the detection of both anti-
HIV-1 and anti-HIV-2 antibodies," J. Immunol. Meth. 139: 77-82 (1991).
148. Srinivasan et al., "Molecular characterization of HIV-1 isolated from a
serum collected in 1976: nucleotide sequence comparison to recent isolates and
generation of hybrid HIV," AIDS Res. Hum. Retrovir. 5: 121-129 (1989).

- 1 149. Starcich et al., "Identification and characterization of conserved and
2 variable regions in the envelope gene of HTLV-III/LAV, the retrovirus of AIDS," Cell
3 45: 637-648 (6/6/86).
- 4 150. Steimer et al., "Neutralization of divergent HIV-1 isolates by
5 conformation-dependent human antibodies to gp120," Science 254: 105-108 (10/91).
- 6 151. Stein & Engleman, "Intracellular processing of the gp160 HIV-1 envelope
7 precursor," J. Biol. Chem. 265: 2640-2649 (2/15/90).
- 8 152. Strebel et al., "A novel gene of HIV-1, *vpu* and its 16-kilodalton
9 product," Science 241: 1221-1223 (9/2/88).
- 10 153. Strebel et al., "Molecular and biochemical analyses of human
11 immunodeficiency virus type 1 Vpu protein," J. Virol. 63: 3784-3791 (9/89).
- 12 154. Streckert & Werchau, "Epitopes at the proteolytic cleavage sites of HIV-
13 1-gp120 and RSV-F protein share a sequence homology: comparative studies with
14 virus-induced and anti-peptide antibodies," Intervirology 34: 30-37 (1992).
- 15 155. Sutcliffe et al., "Antibodies that react with predetermined sites on
16 proteins," Science 219: 660-666 (1983).
- 17 156. Terwilliger et al., "Functional role of human immunodeficiency virus
18 type 1 Vpu," Proc. Natl. Acad. Sci. USA 86: 5163-5167 (7/89).
- 19 157. Thali et al., "Characterization of a discontinuous human
20 immunodeficiency virus type 1 gp120 epitope recognized by a broadly reactive
21 neutralizing human monoclonal antibody," J. Virol. 65: 6188-6193 (11/91).
- 22 158. Thali et al., "Discontinuous, conserved neutralization epitopes
23 overlapping the CD4-binding region of human immunodeficiency virus type 1 gp120
24 envelope glycoprotein," J. Virol. 66: 5635-5641 (9/92).
- 25 159. Thali et al., "Characterization of conserved human immunodeficiency
26 virus type 1 gp120 neutralization epitopes exposed upon gp120-CD4 binding," J. Virol.
27 67: 3978-3988 (7/93).
- 28 160. Thali et al., "Resistance to neutralization by broadly reactive antibodies to
29 the human immunodeficiency virus type 1 gp120 glycoprotein conferred by a gp41
30 amino acid change," J. Virol. 68: 674-680 (2/94).
- 31 161. Veronese et al., "Characterization of gp41 as the transmembrane protein
32 coded by the HTLV-III/LAV envelope gene," Science 229: 1402-1405 (9/27/85).
- 33 162. Vincent et al., "Human immunodeficiency virus type 1 Vpu protein
34 induces degradation of chimeric envelope glycoproteins bearing the cytoplasmic and
35 anchor domains of CD4: role of the Cytoplasmic domain in Vpu-induced degradation
36 in the endoplasmic reticulum," J. Virol. 67: 5538-5549 (9/93).
- 37 163. Wain-Hobson et al., "Nucleotide sequence of the AIDS virus, LAV," Cell
38 40: 9-17 (1/85).

- 1 164. Wain-Hobson et al., "Human immunodeficiency virus type 1 quasispecies
2 in vivo and ex vivo," Curr. Topics Micro. Immunol. 176: 181-193 (1992).
- 3 165. Wang et al., "Detection of antibodies to human T-lymphotropic virus type
4 III by using a synthetic peptide of 21 amino acid residues corresponding to a highly
5 antigenic segment of gp41 envelope protein," Proc. Natl. Acad. Sci. USA 83: 6159-
6 6163 (8/86).
- 7 166. Ward et al., "Decay-accelerating factor CD55 is identified as the receptor
8 for echovirus 7 using CELICS, a rapid immuno-focal cloning method," EMBO J. 13:
9 5070-5074 (11/1/94).
- 10 167. Wells & Compans, "Expression and characterization of a functional
11 human immunodeficiency virus envelope glycoprotein in insect cells," Virology 176:
12 575-586 (1990).
- 13 168. Wiley et al., "Structural identification of the antibody-binding sites of
14 Hong Kong influenza haemagglutinin and their involvement in antigenic variation,"
15 Nature 289: 373-378 (1/29/81).
- 16 169. Willey et al., "In vitro mutagenesis identifies a region within the envelope
17 gene of the human immunodeficiency virus that is critical for infectivity," J. Virol. 62:
18 139-147 (1/88).
- 19 170. Willey et al., "Mutations within the human immunodeficiency virus type 1
20 gp160 envelope glycoprotein alter its intracellular transport and processing," Virology
21 184: 319-329 (1991).
- 22 171. Willey et al., "Human immunodeficiency virus type 1 Vpu protein
23 regulates the formation of intracellular gp160-CD4 complexes," J. Virol. 66: 226-234
24 (1/92).
- 25 172. Willey et al., "Human immunodeficiency virus type 1 Vpu protein
26 induces rapid degradation of CD4," J. Virol. 66: 7193-7200 (12/92).
- 27 173. Willey et al., "Sequences present in the cytoplasmic domain of CD4 are
28 necessary and sufficient to confer sensitivity to the human immunodeficiency virus type
1 Vpu protein," J. Virol. 68: 1207-1212 (2/94).
- 2 174. Wills et al., "Mutations of the rous sarcoma virus Env gene that affect the
3 transport and subcellular location of the glycoprotein products," J. Cell Biol. 99: 2011-
4 2023 (12/84).
- 5 175. Wyatt et al., "Relationship of the human immunodeficiency virus type 1
6 gp120 third variable loop to a component of the CD4 binding site in the fourth
7 conserved region," J. Virol. 66: 6997-7004 (12/92).
- 8 176. Wyatt et al., "Functional and immunologic characterization of human
9 immunodeficiency virus type 1 envelope glycoproteins containing deletions of the
10 major variable regions," J. Virol. 67: 4557-4565 (8/93).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

177. Zhu et al., "Genotypic and phenotypic characterization of HIV-1 in patients with primary infection," Science 261: 1179-1181 (8/27/93)

Date: 5/1/95


B. Matija Peterlin, M.D.

CURRICULUM VITAE

NAME: BORIS MATIJA PETERLIN, M.D.

OFFICE ADDRESS: Rheumatology/Immunology - U426, University of California,
San Francisco, CA 94143-0724

HOME ADDRESS: 14 Hill Point Ave., San Francisco, CA 94117

OFFICE PHONE: (415) 476-1291 and (415) 476-1860

OFFICE FAX: (415) 566-4969 and (415) 476-2956

HOME PHONE: (415) 665-2071

DATE OF BIRTH: July 4, 1947

PLACE OF BIRTH: Ljubljana, Slovenia

CITIZENSHIP: U.S.A.

MARITAL STATUS: Married, 2 children

SSN: 243-62-4412

EDUCATION:

1964 - 1968 Duke University, Durham, North Carolina - B.S. in
Chemistry-Physics, with honors

1968 - 1973 Harvard Medical School, Boston, Massachusetts - M.D.

BOARD CERTIFICATION:

1979 - Diplomate, American Board of Internal Medicine
1980 - Diplomate, American Board of Internal Medicine, Subspecialty Rheumatology

MEDICAL LICENSURE:

State of California - 1973 - present

POSITIONS HELD:

1967 - 1968 NSF Fellowship, Duke University, Durham, North Carolina

1970 - 1971 Teaching Assistant, Harvard Medical School, Harvard
University Research Fellowship

1972 Harvard School of Public Health Fellowship, Apartado,
Colombia

1972 - 1973	CoSTEP
1973	INSERM Fellowship, Institut d'Immunobiologie, Paris, France
1973 - 1975	Intern, Assistant Resident, Medicine, Stanford University Medical Center, Stanford, California
1975 - 1976	Instructor, Georgetown University Medical School, Washington, D.C.
1975 - 1977	Research Associate, LMG, NICHD, NIH, Bethesda, Maryland Lt Cmdr., USPHS
1977 - 1978	Senior Resident, Medicine, Stanford University Medical Center, Stanford, California
1978 - 1980	Leukemia Society Fellow, Clinical Fellow, Division of Immunology, Stanford University Medical Center, Stanford, California
1980 - 1981	American Cancer Society Senior Fellow, Division of Immunology, Stanford University Medical Center, Stanford, California
1981 - 1989	Assistant Professor, Medicine, Microbiology, and Immunology, University of California, San Francisco, California
1981 - 1982	Rosalind Russell Arthritis Scholar, University of California, San Francisco, California
1982 - 1985	Associate, Howard Hughes Medical Institute, University of California, San Francisco, California
1985 - 1991	Assistant Investigator, Howard Hughes Medical Institute, University of California, San Francisco, California
1989 - 1994	Associate Professor, Medicine, Microbiology, and Immunology, University of California, San Francisco, California
1991 - 1995	Associate Investigator, Howard Hughes Medical Institute, University of California, San Francisco, California
1994	Professor of Medicine, Microbiology, and Immunology, University of California, San Francisco, California
1995	Investigator, Howard Hughes Medical Institute, University of California, San Francisco, California

SOCIETIES:

American Association for the Advancement of Science
American Association of Immunologists
American College of Rheumatology
American Federation for Clinical Research
American Society for Clinical Investigation
American Society for Microbiology
International AIDS Society
Phi Beta Kappa
Phi Lambda Upsilon

COMMITTEES:

AIDS Clinical Research Center Committee
American Cancer Society, Member, Study Section on Nucleic Acids and Protein Synthesis
American Cancer Society Scientific Advisory Committee on Molecular Biology and Genetics
Associate Editor, AIDS Res. and Human Retroviruses
Associate Editor, Journal of Immunology
Associate Editor, Virology
Coordinator, Immunology Seminar Series, 1987-1988, 1991-1992
Executive Committee, Program in Biological Sciences (PIBS), 1988-1991
MRC Review of the Genome Studies Section, Member, 1994-
Scientific Committee of Xth International Conference on AIDS, Yokohama, Japan, 1993-1994
Steering Committee, Graduate Program in Immunology, 1988-1991
Transfusion Medicine PPG Internal Advisory Committee
UCSF Dean's AIDS Advisory Committee

BIBLIOGRAPHY

1. Peterlin, B.M., Sullivan, M., Wetphal, H. and Maizel, J.V.: Secondary structure of psoralen cross-linked adenovirus DNA studied by electron microscopy. *Virology* 86:391-397, 1978.
2. Tilghman, S.M., Tiemeyer, D.C., Seidman, J.G., Peterlin, B.M., Sullivan, M., Maizel, J.V. and Leder, P.: Intervening sequence of DNA identified in the structural portion of mouse α -globin gene. *Proc. Natl. Acad. Sci. USA* 75:725-729, 1978.
3. Nossal, N.G. and Peterlin, B.M.: DNA replication by bacteriophage T4 proteins. *J. Biol. Chem.* 254:6032-6037, 1979.
4. Obata, M., Yamawaki-Kataoka, Y., Takahashi, N., Kataoka, T., Simizu, A., Mano, Y., Seidman, J.G., Peterlin, B.M., Leder, P. and Honjo, T.: Immunoglobulin γ 1 heavy chain gene: Structural gene sequence cloned in a bacterial plasmid. *Gene* 9:87-97, 1980.
5. Das, H.K., Biro, P.A., Cohen, S.N., Erlich, H.A., von Gabain, A., Lawrence, S.K., Lemaux, P.G., McDavitt, H.O., Peterlin, B.M., Schulz, M.F., Snod, A.K. and Weissman, S.M.: Use of synthetic probes complementary to human HLA-DR α and β as expression primers for the isolation of S-specific genomic clones. *Proc. Natl. Acad. Sci. USA* 83:1531-1535, 1983.
6. Conwa, T.A., Peterlin, B.M. and Stobo, J.D.: Human I ϵ genes: Structure and function. *Advances in Immunology* 34:71-96, 1983.
7. Payne, R., Brodsky, F.M., Peterlin, B.M. and Young, L.M.: "Bare lymphocytes" in the absence of immunodeficiency. *J. Human Immunology* 6:219-227, 1983.
8. Peterlin, B.M., Conwa, T.A. and Stobo, J.D.: Expression of HLA-DR by a human monocyte cell line is under transcriptional control. *J. Mol. Cell. Immunol.* 1:191-200, 1984.
9. Hardy, K.J., Peterlin, B.M., Atchison, R.E. and Stobo, J.D.: Regulation of expression of the human interferon γ gene. *Proc. Natl. Acad. Sci. USA* 82:8173-8177, 1985.
10. Sullivan, K.E., Stobo, J.D. and Peterlin, B.M.: Molecular analysis of the Bare Lymphocyte Syndrome. *J. Clin. Invest.* 76:73-79, 1985.
11. Calman, A.F. and Peterlin, B.M.: Expression of T cell receptor genes in human B cells. *J. Exp. Med.* 168:1940-1957, 1986.
12. Currey, K.M., Peterlin, B.M. and Maizel, J.V.: Secondary structure of poliovirus RNA: Correlation of computer-predicted with electron microscopically observed structure. *Virology* 148:33-46, 1986.
13. Luciw, P.A., Peterlin, B.M. and Srinivasan, A.: Transactivation of the human immunodeficiency virus. In *Retroviruses of Human A.I.D.S and Related Animal Diseases*. M. Girard, G. De The, L. Valette (Ed), Fondation Marcel Merieux, Lyon, France, pp 61-63, 1986.
14. Peterlin, B.M., Luciw, P.A., Barr, P.J. and Walker, M.D.: Elevated levels of mRNA can account for the transactivation of human immunodeficiency virus. *Proc. Natl. Acad. Sci. USA* 83:9734-9738, 1986.

15. Wang, Y. and Peterlin, B.M.: Methylation patterns of HLA-DRA genes in six mononuclear cell lines. *Immunogenetics* 124:298-303, 1986.
16. Calman, A.F. and Peterlin, B.M.: Mutant human B cell lines deficient in Class II MHC transcription. *J. Immunol.* 139:2489-2495, 1987.
17. Kao, S.Y., Calman, A.F., Luciw, P.A. and Peterlin, B.M.: Anti-termination of transcription within the long terminal repeat of HIV-1 by tat gene product. *Nature* 330:489-493, 1987.
18. Luciw, P.A. and Peterlin, B.M.: Regulated expression of HIV-1 in CD4+ lymphoid cells. In *Retroviruses of Human A.I.D.S. and Related Animal Diseases*. M. Girard, L. Valette. (Eds), Fondation Marcel Merieux, Lyon, France, pp 62-66, 1987.
19. Peltz, G.A., Gallis, B. and Peterlin, B.M.: Monoclonal antibody immunoprecipitation of cell membrane glycoproteins. *Anal. Biol.* 167:239-244, 1987.
20. Peterlin, B.M., Calman, A.F., Luciw, P.A., Salby, M. and Kao, S.-Y.: The mechanism of HIV-1 trans-activation. In *Retroviruses of Human A.I.D.S. and Related Animal Diseases*. M. Girard, L. Valette (Ed), Fondation Marcel Merieux, Lyon, France, pp 83-89, 1987.
21. Peterlin, B.M., Hardy, K.J. and Larsen, A.S.: Chromatin structure of the human HLA-DRA gene in different functional states of MHC class II gene expression. *Mol. Cell Biol.* 7:1967-1972, 1987.
22. Srinivasan, A., Peterlin, B.M., Barr, P.J., Walker, M.D., Rando, R.F., Pellet, P. and Luciw, P.A.: Transactivations of gene expression directed by the long terminal repeat (LTR) of the human immunodeficiency virus (HIV). *Vaccines 87*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, pp 125-129, 1987.
23. Sullivan, K.E. and Peterlin, B.M.: Transcriptional enhancers in the HLA-DQ subregion. *Mol. Cell Biol.* 7:3315-3319, 1987.
24. Sullivan, K.E., Calman, A.F., Nakanishi, M., Tsang, S.Y., Wang, Y. and Peterlin, B.M.: A model for the transcriptional regulation of MHC Class II genes. *Immunology Today* 8:289-293, 1987.
25. Tong-Starko, S.E., Luciw, P.A. and Peterlin, B.M.: Human immunodeficiency virus long terminal repeat responds to T cell activation signals. *Proc. Natl. Acad. Sci. USA* 84:6845-6849, 1987.
26. Wang, Y., Larsen, A.S. and Peterlin, B.M.: A tissue specific enhancer is found in the body of the HLA-DRA gene. *J. Exp. Med.* 166:625-636, 1987.
27. Calman, A.F. and Peterlin, B.M.: Genetic and molecular analysis of TF-XI, a trans-acting factor specific for class II major histocompatibility complex genes. *Proc. Natl. Acad. Sci. USA* 85:8830-8834, 1988.
28. Calman, A.F., Busch, M.P., Vyas, G.N., McHugh, T.M., Sites, D.P. and Peterlin, B.M.: Transcription and replication of human immunodeficiency virus-1 in B lymphocytes *in vitro*. *AIDS* 2:185-193, 1988.
29. Fronck, Z., Timmerman, L.A., Alper, C.A., Hahn, B.H., Kalunian, K., Peterlin, B.M. and McDavitt, H.O.: Major histocompatibility complex associations with SLE. *Am. J. Med.*, 85:542-544, 1988.

30. Luciw, P.A., Tong-Starksen, S.E. and Peterlin, B.M.: Expression directed by the long terminal repeat of the human immunodeficiency virus is enhanced in activated T cells. In *Human Retroviruses, Cancer and AIDS*, D. Bolongue (Ed), UCLA Symposia in Molecular and Cellular Biology, Alan R. Liss, NY, 71:113-122, 1988.
31. Peltz, G.A., Frederick, K., Anderson, C.L. and Peterlin, B.M.: Characterization of the high affinity human monocyte Fc receptor. *Mol. Immunol.* 25:243-250, 1988.
32. Peterlin, B.M.: Genetic aspects of gastro-intestinal immunology. In *Gastrointestinal Immunology*, M. Heyworth, R. Owen, A. Jones (Ed), Raven Press, pp 145-157, 1988.
33. Peterlin, B.M.: Transcriptional regulation of HIV. In *Retroviruses of Human A.I.D.S. and Related Animal Diseases*, M. Girard, L. Valette (Ed), Fondation Marcel Merieux, Lyon, France, pp 95-99, 1988.
34. Peterlin, B.M. and Luciw, P.A.: The molecular biology of HIV. *AIDS* 2:529-540, 1988.
35. Peterlin, B.M. and Luciw, P.A.: Replication of the human immunodeficiency virus: Strategies for inhibition. *Biotechnology* 6:794-799, 1988.
36. Peterlin, B.M., Calman, A.F., Kao, S.-Y., Selby, M.J., Tong-Starksen, S.E. and Luciw, P.A.: Activation and trans-activation of human immunodeficiency virus type 1. *The Control of Human Retrovirus Gene Expression*, B.R. Franza, Jr., B.R. Cullen, F. Wong-Staal (Ed), Cold Spring Harbor Laboratory, pp 45-57, 1988.
37. Peterlin, B.M., Calman, A.F., Kao, S.-Y., Selby, M.J., Tong-Starksen, S.E. and Luciw, P.A.: Transcriptional regulation of human immunodeficiency virus type 1. *Vaccines* 88. H. Ginsberg, F. Brown, R.A. Lerner, R.M. Chanock (Ed), Cold Spring Harbor Laboratory, pp 283-290, 1988.
38. Seto, E., Yen, T.S.B., Peterlin, B.M. and Ou, J.-H.: Trans-activation of the HIV long terminal repeat by the hepatitis B virus X protein. *Proc. Natl. Acad. Sci. USA* 85:8286-8290, 1988.
39. Tsang, S.Y., Nakanishi, M. and Peterlin, B.M.: Tissue specific and interferon- γ inducible regulation of the HLA-DRA promoter. *Proc. Natl. Acad. Sci. USA* 85:8598-8602, 1988.
40. Bain, E.S., Lim, V.L. and Peterlin, B.M.: Designing therapies for the treatment of AIDS. *J. Biotech.* 11:313-324, 1989.
41. Calman, A.F. and Peterlin, B.M.: Transcriptional regulation of HLA class II genes: Cis-acting elements and trans-acting factors. In *Molecular Biology of HLA Class II Antigens*, J. Silver (Ed), CRC Press, pp 225-246, 1989.
42. Peterlin, B.M.: The mechanism of HIV trans-activation by the tat gene product. In *DNA-Protein Interactions in Transcription*, J. Gralla (Ed), UCLA Symposia on Molecular and Cellular Biology New Series 95:221-228, Alan R. Liss, New York, NY, 1989.
43. Selby, M.J., Bain, E., Luciw, P.A. and Peterlin, B.M.: Structure, sequence, and position of the stem-loop in TAR determine transcriptional elongation by Tat through the HIV-1 LTR. *Genes Dev.* 3:547-558, 1989.

44. Seto, E., Zhou, D.-X., Peterlin, B.M. and Yen, T.S.B.: *Trans-activation by the Hepatitis B virus X protein shows cell-type specificity. Virology* 173:764-766, 1989.
45. Tong-Starksen, S.E., Lucht, P.A. and Peterlin, B.M.: Signaling through T lymphocyte surface proteins activates the HIV-1 long terminal repeat. *J. Immunol.* 142:702-707, 1989.
46. Andersson, G.A. and Peterlin, B.M.: NF-X2 that binds to the DRA X2-box is activator protein 1: Expression cloning of c-Jun. *J. Immunol.*, 145:3456-3462, 1990.
47. Barry, P.A., Pratt-Lowe, E., Peterlin, B.M. and Lucht, P.A.: Cytomegalovirus activates transcription directed by the long terminal repeat of the human immunodeficiency virus type 1. *J. Virol.* 64:2932-2940, 1990.
48. Fronck, Z., Tlunerman, L.M., Alper, C.A., Hahn, B.H., Kalunian, K., Peterlin, B.M. and McDewitt H.O.: Major histocompatibility complex genes and susceptibility to systemic lupus erythematosus. *Arth. Rheum.* 33:1542-1553, 1990.
49. Peterlin, B.M.: Regulation of HIV gene expression. *Human retroviruses*, Croopman, J.E., Chen, I., Essex, M. and Weiss, R. (Eds), *UCLA Symposia on Molecular and Cellular Biology* 119:153-162, Alan R. Liss, New York, NY, 1990.
50. Peterlin, B.M., Andersson, G.A., Lotscher, E. and Tsang, S.Y.: Transcriptional regulation of HLA class II genes. *Immunol. Res.* 9:164-177, 1990.
51. Selby, M.J. and Peterlin, B.M.: *Trans-activation by HIV-1 Tat via a heterologous RNA-binding protein. Cell* 63:769-776, 1990.
52. Tong-Starksen, S.E., and Peterlin, B.M.: Differences in transcriptional enhancers of HIV-1 and HIV-2: Response to T cell activation signals. *J. Immunol.* 145:4348-4354, 1990.
53. Tong-Starksen, S.E. and Peterlin, B.M.: Mechanisms of retrovirus transcriptional activation. *Seminars in Virology* 1:215-227, 1990.
54. Tsang, S.Y., Nakanishi, M. and Peterlin, B.M.: Mutational analysis of the DRA promoter: cis-acting sequences and trans-acting factors. *Mol. Cell Biol.* 10:711-719, 1990.
55. Chin, D.J., Selby, M.J. and Peterlin, B.M.: HIV-1 Tat does not trans-activate mature TAR RNA species in the nucleus or cytoplasm of primate cells. *J. Virol.* 65:1758-1764, 1991.
56. Derse, D., Carvalho, M., Carroll, R. and Peterlin, B.M.: A minimal lentivirus Tat. *J. Virol.* 65:8333-8337, 1991.
57. Modesti, N., Garcia, J., Debouck, C., Peterlin, B.M. and Gaynor, R.: Trans-dominant mutants resulting alterations in the Tat basic domain inhibit HIV gene expression. *New Biologist* 3: 759-760, 1991.
58. Peterlin, B.M.: Transcriptional regulation of HIV. *Genetic structure and regulation of HIV*, W.A. Haseltine, F. Wong-Staal (Ed.), Raven Press, Boca Raton, FL, pp. 237-250, 1991.
59. Peterlin, B.M.: Transcriptional regulation of HLA-DRA gene. *Immunol. Res.* 142:393-399, 1991.

60. Selby, M.J. and Peterlin, B.M.: HIV-1 Tat trans-activates in the absence of its target. *Advances in Molecular Biology and Targeted Treatment of AIDS*, A. Kumar (Ed), Plenum Press, New York, NY, pp. 123-132, 1991.
61. Alonso, A., Derse, D. and Peterlin, B.M.: Human chromosome 12 is required for optimal interactions between Tat and TAR of HIV-1 in rodent cells. *J. Virol.* 66:4617-4621, 1992.
62. Carroll, R., Peterlin, B.M. and Derse, D.: Inhibition of HIV-1 Tat activity by co-expression of heterologous trans-activators. *J. Virol.* 66:2000-2007, 1992.
63. Voliva, C.F., Amrhein, A., Walker, M.D. and Peterlin, B.M.: B-cell factor 1 is required for optimal expression of the DRA promoter in B cells. *Mol. Cell. Biol.* 12:2383-2390, 1992.
64. Ghosh, S., Selby, M.J. and Peterlin, B.M.: Transcriptional synergy between Tat and VP16 on the HIV-1 LTR. *J. Mol. Biol.* 234:610-619, 1993.
65. Lu, X., Welsh, T., and Peterlin, B.M.: The human immunodeficiency virus type 1 long terminal repeat specifies two different transcription complexes, only one of which is regulated by Tat. *J. Virol.* 67: 1752-1760, 1993.
66. Luo, Y. and Peterlin, B.M.: Juxtaposition between activation and basic domains of HIV-1 Tat is required for optimal interactions between Tat and TAR. *J. Virol.* 67:3441-3445, 1993.
67. Luo Y., Madore, S.J., Parslow, T.G., Cullen, B.R. and Peterlin, B.M.: Functional analysis of interactions between Tat and TAR of HIV-1 in cells. *J. Virol.* 67:5617-5622, 1993.
68. Peterlin, B.M.: Introduction: Transcriptional Regulation of Viruses. *Seminars in Virology*, 4:1, 1993.
69. Peterlin, B.M. (Ed), Transcriptional Regulation of Viruses. *Seminars in Virology*, 4:1-80, 1993.
70. Peterlin, B.M.: Intracellular factors influencing HIV replication: Discussion. In *Focus on HIV*, Neu, H., Levy, J.A., and Weiss, R.A. (Eds), Churchill Livingstone, Edinburgh, United Kingdom, pp 96-97, 1993.
71. Peterlin, B.M., Adams, M., Alonso, A., Baur, A., Ghosh, S., Lu, X. and Luo, Y.: Tat trans-activator. in *Molecular Biology of Human Retroviruses*. Cullen, B. R. (Ed), IRL/Oxford University Press, Oxford, United Kingdom, pp 75-100, 1993.
72. Tong, S.B., Lu, X., Baur, A. and Peterlin, B.M.: Second exon of Tat of HIV-2 is required for optimal trans-activation of HIV-1 and HIV-2 LTRs. *Virology* 199:826-830, 1993.
73. Voliva, C.F., Tsang, S.Y. and Peterlin, B.M.: Mapping cis-acting defects in promoters of transcriptionally silent DQA2, DQB2, and DOB genes. *Proc. Natl. Acad. Sci. USA* 90:3408-3412, 1993.
74. Zhang, X.-Y., Jabrane-Ferrat, N., Astuda, C.K., Peterlin, B.M. and Ehrlich, M.: The major histocompatibility complex class II promoter-binding protein RFX (NF-X) is a methylated DNA-binding protein. *Mol. Cell. Biol.* 13:6810-6818, 1993.
75. Adams, M., Sharmeen, L., Kimpton, J., Romeo, J.M., Garcia, J.V., Peterlin, B.M., Groudine, M. and Emerman, M.: Cellular latency in human immunodeficiency virus-infected

individuals with high CD4 levels can be detected by the presence of promoter-proximal transcripts. *Proc. Natl. Acad. Sci. USA* 91:3862-3866, 1994.

76. Alonso, A., Cujec, T. and Peterlin, B.M.: Effects of human chromosome 12 on interactions between Tat and TAR of human immunodeficiency virus type 1. *J. Virol.* in press, 1994.
77. Baur, A. and Peterlin, B.M.: Effects of CD45 on NF- κ B: implications for replication of HIV-1. *J. Immunol.* 152:976-983, 1994.
78. Baur, A.S., Sawai E.T., Dazin, P., Fandl, W.J., Cheng-Mayer, C. and Peterlin, B.M.: HIV-1 Nef leads to inhibition or activation of T cells depending on its intracellular localization. *Immunity*, 1:373-384, 1994.
79. Chang, C.-H., Fontes, J.D., Peterlin, B.M., and Flavell, R.A.: CIITA is sufficient for the inducible expression of class II genes. *J. Exp. Med.* 180: 1367-1374, 1994.
80. Jabrane-Ferrat, N., and Peterlin, B.M.: Ets-1 activates DRA promoter in B cells. *Mol. Cell. Biol.* 14: 7314-7321, 1994.
81. Jones, K.A. and Peterlin, B.M.: Control of RNA initiation and elongation at the HIV-1 promoter. *Annu. Rev. Biochem.* 63:717-743, 1994.
82. Luo, Y. and Peterlin, B.M.: RNA-binding trans-activators in cells. In *Molecular Virology Techniques*, Adolph, K.W. (Ed), Academic Press, New York, NY, pp29-43, 1994.
83. Luo, Y., Yu, H. and Peterlin, B.M.: Cellular protein modulates effects of human immunodeficiency virus type 1 Rev. *J. Virol.* 68:3850-3856, 1994.
84. Mujeeb, A., Bishop, K., Peterlin, B.M., Turck, C., Parslow, T.G. and James, T.L.: NMR structure of a biologically active peptide containing the RNA-binding domain of HIV-1 Tat. *Proc. Natl. Acad. Sci. USA* 91:8248-8252, 1994.
85. Sawai, E.T., Baur, A.S., Struble, H., Peterlin, B.M., Levy, J.A. and Cheng-Mayer, C.: HIV-1SF2 Nef associates with a cellular protein kinase. *Proc. Natl. Acad. Sci. USA*, 91: 1539-1543, 1994.
86. Wright, S., Lu, X., and Peterlin, B.M.: Human immunodeficiency virus type 1 Tat direct transcription through attenuation sites within the mouse c-myc gene. *J. Mol. Biol.* 243: 568-573, 1994.
87. Itoh-Lindstrom, Y., Peterlin, B.M. and Ting, J.P.-Y.: Affinity enrichment and functional characterization of TRAX1, a novel transcription activator and X1 sequence binding protein of HLA-DRA. *Mol. Cell. Biol.* 15:282-289, 1995.
88. Peterlin, B.M.: Molecular Biology of HIV. In *Retroviridae*, Levy, J.A. Ed., Plenum Press, New York, NY, in press, 1995.
89. Sawai, E.T., Baur, A.S., Peterlin, B.M., Levy, J.A., Cheng-Mayer, C.: A highly conserved region and membrane targeting of Nef from primate immunodeficiency viruses are required for association with a cellular serine kinase activity. *J. Biol. Chem.* in press, 1995.

90. Haldsinger, A.A., Abshire, K.Z., Peterlin, B.M. and Antonucci, T.: HIV-1 Nef repression of HIV-1 LTR driven transcription is independent of protein kinase C activity in Jurkat cells. *Virology* submitted, 1995.
91. Voliva, C.F., Jabrane-Ferrat, N. and Peterlin, B.M.: The octamer-binding site participates in TFIID recruitment to the DRA promoter. *J. Immunol.* submitted, 1995.
92. Jabrane-Ferrat, N., Nakanishi, M. and Peterlin, B.M.: RF-X binds to the S box in the DRA promoter. *Mol. Cell. Biol.* submitted, 1995.

THIS PAGE BLANK (USPTO)